

PROCESSES FOR PRODUCTION OF IMMUNOGLOBULIN A IN MILK

5

TECHNICAL FIELD

10 This invention relates to processes for producing immunoglobulin A in mammals, processes for producing milk containing immunoglobulin A and to the uses of the immunoglobulin A and milk produced.

BACKGROUND

- 15 Immunoglobulin A (IgA) is a well documented immunoglobulin present in almost all body fluids. It is thought to play a major role in the protection of the host from infection by pathogenic organisms invading via the mucosal surfaces of the respiratory, gastrointestinal and urogenital tracts. IgA participates in the clearance of pathogenic bacterial, viral or parasitical organisms and a variety of ingested or inhaled antigens from the mucosal surfaces by neutralising toxins and viral particles, inhibiting adherence of
- 20 bacterial pathogens and preventing colonisation and penetration of mucosal surfaces by pathogenic microorganisms. The key role of immunoglobulins including IgA in milk therefore is to provide local protective immunity in the gastrointestinal tract of the offspring during the suckling period.
- 25 Immunoglobulins have come to be recognised as useful in the pharmaceutical and veterinary fields for treating bacterial or viral infections of the gut, and more generally in the treatment of disease and inflammation. Over the years various techniques for producing immunoglobulins have been proposed. A particularly popular method is for the induction and harvesting of immunoglobulins from ruminant milk. This approach has
- 30 particular advantages in that the immunoglobulin produced in the milk is in a form suitable for immediate consumption, or may be processed into appropriate formulae or products. It is safe to use and the industry infrastructure for producing milk containing antibodies is already in place.
- 35 The ruminant immune system appears to differ from its human counterpart in that the immunoglobulin dominant in bovine mammary secretions is IgG₁. Accordingly, the main focus of antibody production in milk by active immunisation has been on immunoglobulin G's, although theoretically, the preferred immunoglobulin would be IgA for the reasons

outlined above.

Some attempts have been made to produce increased levels of IgA in ruminant milk. Proposals for vaccination by a single administration route such as parenteral, subcutaneous, intravenous, systemic, oral, intraperitoneal, intramuscular, intramammary and the like have been suggested. Generally, these routes of administration have resulted in the predominant production of IgG₁. Systemic immunisation produced both IgA and IgM in milk, but only at low concentrations. The response was enhanced when intramuscular/sub-cutaneous (IM) and intramammary (IMM) immunisation processes were combined (Am. J. Vet. Res¹). Combinations of intraperitoneal (IP) and intramammary (IMM) infusion have also been shown to produce IgA and IgG₁ (Immunology⁷; Res. in Vet.Sci⁸, Res. in Vet.Sci¹¹, The Ruminant Immune System in Health and Disease¹⁰). It is noted that this route leads to limited enhancement of IgA production (The Ruminant Immune System in Health and Disease¹⁰). A combination of IM and IMM immunisation gave rise to a predominance of IgG₁ in the milk (Aus. J. Dairy Technology⁶), as well as increasing generally the levels of IgG₂, IgA, and IgM (Am. J. Vet. Res¹). Significant between animal variability in the antibody titres produced was also noted.

The predominance of the production of IgG₁ is consistent with the findings that IgG's produced are the major immunoglobulins in ruminant mammary secretions.

Intramammary immunisation techniques have generally not been preferred as a route for vaccination under field conditions due to the high chance of mammary infection (Aus. J. Dairy Technology⁶). However, other work suggests that this may not be the case (Am. J. Vet. Res¹).

It should be noted that much of the published literature concerning immunoglobulin production in mammary gland secretions is directed to disease prevention in animals or their offspring. Few are directed to the production of immunoglobulin enriched milk for the purposes of obtaining the immunoglobulins themselves.

An exception to this is a process for the production of a protein concentrate containing immunological factors of lactic origin in Swiss Patent No. 1,573,995. Nearly 20 years ago, this patent disclosed a process for producing milk with a high antibody titre, by intracisternal instillation into the mammary gland, parental injection (subcutaneous, intravenous), injection into the retromammary ganglionic system by scarification, by oral ingestion or by a combination of several of these modes. The only specific immunisation

protocol for obtaining colostral and transition milk disclosed involved some 11 immunisation steps over a period of 8 weeks prior to calving. This protocol comprises multiple parental (including intravenous) administration steps, with several IMM administration steps interspersed and requires 2 oral administration steps in the week prior to calving.

This protocol is not in widespread use today. The immunisation plan is onerous in the number of steps involved and is not in fact optimised for immunoglobulin A production. Indeed, the patent is misleading in suggesting that IgA's are preponderant in ruminant maternal milk; a misconception that may have resulted from the knowledge that IgA is predominant in human milk. As established in other teachings (see for example, Aus. J. Dairy Technology⁶), IgG is the predominant immunoglobulin produced in the maternal milk.

It has also been shown in the intervening years that oral delivery of antigens results in little or no increase of IgA titres in mammary secretions when compared with non-inoculated controls (Am. J. Vet. Res¹). It is presumed that the presence of the rumen may preclude the antigen reaching the small intestine. Accordingly, the oral administration step called for by Hilpert is now contraindicated.

Similarly, intravenous injection would not generally be recommended for immunisation purposes because of the possible adverse effects such as anaphylactic shock (Cold Spring Harbour¹⁸, ILAR Journal¹⁹).

There is currently a need for a process for inducing, and producing IgA in milk at higher levels than have previously been obtained by known antigen administration processes. A process which additionally reduces between animal variability in the production of IgA is also desirable. A commercial process which optimises production of IgA while simplifying the immunisation protocol is also sought.

It is therefore an object of this invention to provide a process for the induction and production of immunoglobulin A in milk which goes some way towards overcoming the above disadvantages or at least provides the public with a useful choice.

Accordingly, the present invention can broadly be said to consist in a process for the induction of immunoglobulin A (IgA) in a mammal which process comprises:

(a) actively immunising a pregnant mammal with an antigen by any two routes of

administration selected from intramammary (IMM), intraperitoneal (IP) and intramuscular (IM); and

- (b) actively immunising said mammal with an antigen by a third administration route selected from intramammary (IMM), intraperitoneal (IP) and intramuscular (IM);
5 with the proviso that all three administration routes are different.

In a further aspect the present invention provides a process for the production of mammalian milk containing immunoglobulin A (IgA), which process comprises:

- 10 (a) induction of IgA according to the process set out above; and
(b) collecting milk containing IgA from said mammal.

Preferably, the initial immunisation protocol is followed by a programme of booster immunisations over the preparturition period.

- 15 In a preferred process of the present invention the antigen administered is the same for each route of administration.

- Preferably, the antigen administered is emulsified in an adjuvant. A particularly preferred
20 adjuvant is Freund's incomplete adjuvant (FIC).

In one embodiment of the invention IgA may be isolated from the mammalian milk collected. The isolated IgA may be purified if desired.

- 25 In a further aspect, the present invention provides mammalian milk containing IgA produced in accordance with the processes of the invention.

In a still further aspect, the present invention provides IgA produced in accordance with the processes of the invention.

- 30 Preferred mammals for use in the processes of the present invention are ruminants, especially dairy cows.

- The present invention further provides for the use of immunoglobulin A produced in
35 accordance with the processes of the invention in pharmaceutical, cosmetic and veterinary compositions as well as in food products including functional foods and dietary supplements.

Although the present invention is broadly as defined above, it will be appreciated by those persons skilled in the art that the invention is not limited thereto and that it also includes embodiments of which the following description gives examples. In particular, preferred aspects of the invention will be described in relation to the accompanying drawings in which:

Figure 1 shows a typical immunoglobulin dilution curve for a positive control sample in the ovine *E.coli* IgA enzyme linked immunoabsorbent assay (ELISA).

- 10 Figure 2 shows the right (immunised) gland milk *anti-E.coli* IgA titres for all groups on day 0, day 5, week 2 and week 4 after parturition.

Figure 3 contrasts *anti-E.coli* IgA milk responses of the right (immunised) gland and the left (untreated) gland on day 0.

- 15 Figure 4 shows the *anti-E.coli* IgA responses for individual sheep with right (immunised) glands on day 1 post-partum. The effect of the immunisation route is depicted.

- 20 Figure 5 shows the *anti-E.coli* responses for individual sheep with right (immunised) glands on day 2 post-partum.

Figure 6 shows the *anti-3K* scour guard right (immunised) gland milk IgA antibody titres for all groups on days 0 and 5.

- 25 Figure 7 shows the *anti-3K* scour guard right (immunised) gland and left gland milk IgA antibody titres for all groups on day 0.

Figure 8 shows a typical dilution curve for positive and negative control samples in the ovine TNF IgA ELISA.

- 30 Figure 9 shows individual sample analyses of *anti-TNF* IgA titre responses in the right (immunised) gland day 1 *post-partum* milk samples for each of the immunisation groups.

- 35 Figure 10 shows the data for *anti-TNF* titres from day 1 post-partum samples from the right (immunised) and left (untreated) gland.

Figure 11 shows the relationship of *anti-TNF* IgA to stage of lactation.

Figure 12 shows a typical dilution curve for positive and negative control samples in the C.albicans IgA ELISA.

Figure 13 shows the right (IMM immunogen in FIC) gland milk anti-C.albicans IgA antibody titre for all groups on Day 1, 2, 7, 14 and 60.

Figure 14 contrasts the anti-C.albicans IgA response in the left gland (aqueous IMM immunogen) and right (FIC IMM immunogen) glands over the lactation period.

10 DETAILED DESCRIPTION

The term "milk" used herein refers to both milk and colostrum in the form in which it is produced by the mammal.

15 The term "antigen" as used herein refers to any material capable of inducing an antigenic response in a treated mammal.

In a first aspect the present invention relates to a process for the induction of immunoglobulin A (IgA) in a mammal. As a first step the method comprises actively immunising a pregnant mammal with an antigen by any two administration routes selected from intramammary (IMM), intraperitoneal (IP) and intramuscular (IM). As a second step the mammal is again actively immunised by a third administration route selected from those routes given above. The proviso to this process is that all three administration routes selected are different.

25 The applicants have surprisingly found that the use of three routes of administration increases IgA antibody titre levels above what might be expected by simply combining two known administration route processes with a third route of administration, or at least decreases the between animal variability in IgA antibody titre response.

30 It will be appreciated by the reader that the ordering of the routes and timing of administration is not crucial to the process for inducing immunoglobulin A. Moreover, the immunisations by the different routes may be carried out sequentially, discontinuously or concurrently. A presently preferred immunisation protocol is for concurrent IM and IP immunisation followed by IMM immunisation. The IM and IP immunisations effectively act to prime the immune system response. The IMM immunisation is a localised challenge to induce IgA production in that immunised region.

In a further preferred embodiment the initial immunisation protocol is followed by a number of booster doses of antigen over the pre-parturition period. The amounts of antigen introduced, the frequency (time interval), and the number of booster doses may vary widely. For example, from a single booster shot via a single route of administration on one occasion through to multiple shots via each of three administration routes on many different occasions.

Booster shots are generally spaced to suit the convenience of the operator. To avoid local irritation and congestion, it is usually preferred that booster shots to the same site not be given more frequently than every other week.

One regimen preferred requires concurrent IM and IP immunisation on two separate occasions, followed by IMM immunisation on one occasion. That is, effectively two priming steps followed by local challenge. The first priming step is generally carried out 2 to 8 weeks before the second priming and challenge steps. These latter steps are desirably carried out concurrently. A convenient protocol is for the first priming step to be effected 6 to 14, preferably 8 to 12, and most desirably, 8 weeks before parturition and the second priming/local challenge step to be effected 2 to 10 weeks, preferably 4 to 8, and most desirably, 4 weeks before parturition. However, as noted above timing is not crucial.

A second preferred regimen is for initial immunisation 6 to 14, preferably 8 to 12 and most desirably, 8 weeks before parturition followed by 1 or 2 booster shots via each of the three administration routes on 1 to 3 occasions before parturition. The final immunisation will generally be given 1 to 2 weeks before parturition.

Particularly preferred is a regimen requiring one additional priming and local challenge step such that at 8 weeks prior to parturition (minus 8 weeks) there is a concurrent IM and IP immunisation, followed by concurrent IM/IP and IMM immunisations at minus 4 weeks, a second IMM immunisation at minus two weeks, and a final concurrent IM and IP immunisation at minus one week.

A further preferred regimen is for the initial IM/IP immunisations to be effected at 12 weeks before parturition and the second priming/local challenge step at 8 weeks before parturition, a second IMM immunisation at minus 6 weeks and a final concurrent IM/IP immunisation at minus 4 weeks.

It will be appreciated from the above that a wide variation in the timing of the

immunisations is feasible generally starting 14 weeks prior to parturition, but preferably 12 or 8 weeks prior to parturition.

Subsequent to parturition, declining antibody concentrations can be increased by periodically introducing booster shots of the selected antigen into the mammal during the lactation period according to equivalent preparturition protocols outlined above. Generally, this involves between 1 to 6, preferably 2 to 4, and most preferably 2 or 3, concurrent IM and IP immunisations in the lactation phase following parturition together with 1 IMM immunisation at the involution stage of lactation.

10

In a further embodiment of the invention the process of the invention further comprises a preselection step. In this step individual animals are tested and selected for their ability to produce IgA.

As noted above, considerable between animal variability exists for the production of immunoglobulins. This preselection step wherein the animals showing the best IgA antibody titre responses are selected assists in decreasing the between animal variability factor. This process may similarly be used to build groups of animals particularly suited to IgA production.

20

Processes for IM, IP and IMM administration are well known in the art. For IM immunisation it is generally preferred that more than one site be used for administration by this process. Preferred sites for IM administration are the left and right sides of the brachio cepalic muscle (that is, two sites in one muscle). For IP immunisation, administration into the peritoneal cavity, generally at only one site is presently preferred. Desirably, administration is at the sub lumbar fossa. The precise sites of administration for these routes may of course vary according to known administration protocols. The amount and form of the antigen administered will also vary according to the antigen used and the mammal to be immunised in accordance with known vaccine formulations.

30

Generally, the antigen is injected using the syringe and needle for IM and IP routes and fine-bore polyethylene surgical tubing fitted to a syringe for the IMM route or alternatively a conventional sterile intra mammary applicator. For the IMM immunisation, the antigen is generally administered via the major lactiferous duct or the supramammary lymph node. Preferably, via the teat orifice into the teat canal. For best results it is also preferred that each mammary gland be immunised on each occasion. This maximises the localised IgA response mounted in the mammal.

35

The volume of antigen injected will vary according to the mammal and the route of immunisation. Table 1 below is a summary of the injection volumes for sheep and cows immunised via the IM, IP and IMM routes.

5

Table 1

Sheep

	IM	IP	IMM
10 volume	1.0 ml (per site)	1.0 ml	1.0 ml (per gland)
maximum volume	5.0 ml	2.5 ml	2.0 ml (per gland)

Cows

	IM	IP	IMM
15 volume	2.0 ml (per site)	4.0 ml	2.0 ml (per gland)
maximum volume	8.0 ml	10.0 ml	5.0 ml (per gland)

20 Typically for immunising the bovine, antigen is administered at 2 ml per site and 2 sites for IM, 4 ml at 1 site for IP and 2 ml into each of the four glands for IMM.

Contrary to conventional wisdom, field trials show that there is no significant risk of infection using intramammary immunisation provided appropriate precautions are taken. For example, care must be taken to sterilise glands prior to immunisation. Appropriate
25 sterilisation methods are known in the art. For example, ethanol/iodine washes will serve this purpose. A further precaution is to ensure that the antigen is administered in a solution containing antibiotic. Suitable antibiotics include dupocillin and ampicillin and clavulox L.C.

30 The mammals selected for use in the process of the invention will generally be economically useful mammals such as ruminants. Examples of the ruminants preferred for use are cows, goats, and sheep.

35 The term "antigen" as used herein refers to any material capable of inducing an antigenic response in the treated mammal. Antigens may be selected according to the ultimate utility of the IgA formulation. That is, if the formulation is to be used for generating passive immunity, the antigen against which such immunity is sought should be used. Antigenic substances which may be employed in the process of the invention include

bacteria, viruses, yeasts, mycoplasmas, proteins, haptens, animal tissue extracts, plant tissue extracts, spermatozoa, fungi, pollens, dust, chemical antigens and mammalian cells.

- Where haptens are to be used as antigens these should first be conjugated to carrier substances such as proteins using chemistry well known to people versed in the art. (ILAR Journal¹⁹).

- Useful bacterial antigens include species of Escherichia, Staphylococcus, Streptococcus, Salmonella and Pneumococcus. Particularly preferred bacterial antigens are
- 10 Escherichia coli, clostridium difficile, vibrio cholerae and helicobacter pylori.

Preferred yeast antigens include species of Candida.

- A particularly preferred yeast antigen is Candida albicans.
- 15

- Useful viral antigens include rotavirus, herpes, fowlpox, rhinopneumonitis, coronavirus, parvovirus and influenza. Protein antigens include tumour necrosis factor, insulin-like growth factors, and somatostatin, viral or bacterial cell surface proteins and conjugated protein antigens. Chemical antigens include pollens, pesticides, insecticides, fungicides
- 20 and toxins. Complex antigens comprising a combination of two or more antigens of the types identified are also feasible. One such preferred complex antigen is 3K Scourguard (SmithKline Beecham, Royal Oak, Auckland, New Zealand). The vaccine contains pathogenic E.coli, bovine rotavirus, and coronavirus.

- 25 Useful mycoplasma antigens include mycoplasma pneumoniae and cryptosporidium parvum.

- Generally, the antigenic substances are suspended in liquid medium for infusion or injection according to known protocols. Any appropriate carriers, diluents, buffers, and
- 30 adjuvants known in the art may be used. Suitable suspension liquids include saline solution, water, and physiologic buffers.

- The use of adjuvants is also desirable. Suitable adjuvants for use with the antigens of the invention include Freund's complete adjuvant (FCA), Freund's incomplete adjuvant
- 35 (FIC), adjuvant 65, cholera toxin B subunit, alhydrogel; or *bordetella pertussis*, muramyl dipeptide, cytokines and saponin. Oil based adjuvants and in particular FCA and FIC are preferred.

Prior to injection antigens in appropriate carriers are typically emulsified with oil-based adjuvant (FIC is preferred) using a laboratory homogeniser. Aqueous antigen is typically mixed with 3 volumes of oil adjuvant and emulsified until a stable water in oil emulsion is formed as demonstrated using tests well known in the art.

- 5
- Conventional wisdom also taught that the use of oil based adjuvants with direct intramammary immunisation was not feasible because of the risk of adverse reactions. The present applicants have found that not only is the administration with oil based adjuvants feasible under appropriate care conditions, but is also desirable. The use of FIC may
- 10 significantly enhance the immunogenic response obtained for some antigens, particularly when administered by the IMM route. It is therefore presently preferred that antigens be emulsified in FIC for all immunisations except for small polypeptides where FCA may be preferable for the first IM/IP immunisation. However, FCA has not been used for IMM.
- 15
- As noted above, the size and concentration of the antigen doses are not critical and it is known in the art that there is a dose range known as the window of immunogenicity for antigen and that this is generally relatively broad.
- 20
- However, too much or too little antigen may induce suppression, tolerance or immune deviation towards cellular immunity and away from humoral immune response. Typically, for protein antigens optimal doses are of the order of 5 to 25 $\mu\text{g/kg}$ live weight in ruminants and for dead, lyophilised bacterial or viral antigens doses in the range of 1×10^8 to 4×10^{10} organisms per ml are typical.
- 25
- As also noted above, in IMM immunisation it is preferred that the antigen additionally be formatted in suspension with an antibiotic.
- 30
- Regarding the specific form of the antigen, it will be appreciated by the reader that both live and killed vaccines are possible. Studies have shown that killed vaccines will stimulate IgG_1 responses while live vaccines stimulate IgG_2 responses. For the production of IgA either alternative is possible.
- 35
- The antigen administered by each of the immunisation routes may be the same or different. Accordingly, several different antigens may be administered by the three different immunisation routes for each of the initial and booster immunisations. However, it is presently preferred that the same antigen or combination of antigens be administered

via the three routes on each immunisation occasion.

5 In a further aspect the present invention relates to a process for the production of mammalian milk containing IgA which method comprises induction of IgA antibodies according to the process detailed above and then collecting the IgA containing milk from the mammal. The collection of milk may be effected using normal milking processes.

10 IgA titre responses are generally highest on the first day following parturition. After this antibody levels drop to between 5 to 20% of the initial level. This subsequent level is usually maintained for two to three months or until drying off or involution. IgA levels may be raised in this period through booster immunisations as discussed above. Milk containing IgA may usefully be collected throughout this period.

15 This milk is useful in the form obtained directly from the mammal but may be processed if required. Examples of processing steps include heat treatment, ultra violet radiation, concentration, supplementation with food additives, drying into concentrates, milk powders and the like.

20 As a further step to the process of the invention, the IgA may be isolated from the milk. Isolation may be effected using separation techniques known in the art. For example, isolation of immunoglobulin rich fractions from whey in Can. J. Vet. Res²¹, EP 0320152, WO 97/27757, GB2179947, from milk in Milchwissenschaft²², US 4,229,342, from colostrum in Agric. Biol. Chem²⁰, French Patent No. 2520235, New Zealand Patent No. 239466 and US 4,582,580, and from milk and colostrum in US 4,644,056.

25 The isolated IgA may subsequently be purified if desired. Purification may be carried out according to known techniques such as precipitation and ion exchange chromatography. Suitable techniques are disclosed in the journals and patents referenced above. Both the isolated and purified immunoglobulin A produced in accordance with the additional
30 process steps also form part of the present invention.

In a further aspect the present invention relates to mammalian milk containing IgA produced in accordance with the process of the invention.

35 Processes for producing protein concentrates containing immunoglobulins on a commercial scale are disclosed in Swiss Patent No. 1,573,995 incorporated herein by reference. Briefly, the process comprises collecting the milk of hyperimmunised milk-bearing females; separating the cream and the impurities, coagulating the clarified and

skimmed milk, separating the casein, filtering, ultrafiltering and sterilising the proteins of the whey by filtration, evaporating and drying the product under conditions which do not denature the immunoglobulins and which maintain sterility.

- 5 In a further aspect, the present invention provides for the use of IgA in the form of milk, processed milk products, concentrates, isolated IgA and purified IgA produced in accordance with the process of the invention. IgA has potentially broad applications in the fields of pharmaceutical, veterinary and cosmetic compositions as well as in foods and dietary supplements. Such compositions, food and supplements can be administered to
10 patients (including human patients) having need of same.

More specifically, in the human health care field passive oral immunisation using milk immunoglobulins from specifically vaccinated cows has long been known. Given the significant role that IgA plays in preventing enteric infections, formulations containing
15 IgA may be effective in treating patients susceptible to such enteric infections. Formulations containing IgA antibodies against enterotoxigenic gastric pathogens including pathogenic *E.coli*, Rotavirus, *Staphylococcus*, *Streptococcus*, *Aerobacter*, *Salmonella*, *Pseudomonas*, *Haemophilus influenza*, *proteus vulgaris*, *shigella dysenteriae*, *Diplococcus pneumoniae*, coronavirus and *Corynebacterium acne* are all possible.

20 Formulations containing high levels of IgA specific for infants is one application. Infants are often very susceptible to enteric gastric disorders. Specific formulations containing anti-cryptosporidiosis IgA for protection against cryptosporidiosis infection in HIV and AIDS patients is a further possibility. General formulations for protection of travellers
25 against diarrhoea and general gastric disorders are contemplated. Valuable formulations containing antibodies against *Helicobacter pylori* for protection against stomach ulcers are feasible.

Appropriate formulations can be produced based on known art formulations. For
30 example, formulations for treating the following disorders are provided for in the art:

	<u>Treatment of</u>	<u>Reference</u>
	Gastroenteritis	Swiss Patent No. 1,573,995
35	Infantile <i>E.coli</i> gastroenteritis	Eur. J. Pediatr. ¹⁴
	Enteric infections	Advances in Exp. Med. & Biol. ¹⁵
	Enteric disease	US 5,066,491
	Campylobacter jejuni	J. Applied Bacteriology ²⁵

	Shigella Flexneri	Am. J. Tropical medicine and Hygiene ²⁷
	Rotavirus diarrhoea	Indigenous Antimicrobial Agents ²³
	Dental Caries	Infection and Immunity ²⁸
	Cryptosporidial diarrhoea	Lancet ²⁴ , Gastroenterology ²⁹
5	Cryptosporidiosis in AIDS	Archives of Disease in Childhood ³⁰
	Rotavirus gastroenteritis	J. Infectious Diseases ¹⁶ , J. Clinical Microbiology ¹⁷
	H.pylori	US 5,260,057
	Respiratory disease	US 5,066,491
10	Cryptosporidiosis	US 5,066,491

15 A comprehensive review of the use of bovine immunoglobulins to treat or prevent certain human diseases caused by H.pylori, C.parvum, E.coli, S.flexneri, C.difficile, V.cholerae and rotavirus is provided in the proceedings of the IDF seminar on Indigenous antimicrobial agents of milk²³.

In a more general context, pharmaceutical formulations containing IgA tailored to the needs of the young, old, medically impaired, and terminally ill are all desirable.

20 The formulations of the invention similarly have applications in the veterinary field. For example in the preparation of formulations containing specific IgA antibodies against pathogenic microbiologies such as E.coli, rotavirus, coronavirus and other scour causing microbes for the prevention and treatment of gastric disorders in neonatal livestock.

25 Formulations containing specific IgA antibodies against mycotoxins, phytotoxins, aflotoxins, herbicides, pesticides and fungicides to block absorption of these following oral ingestion are possible.

30 More generally, formulations may be prepared containing IgA against undesirable food ingredients to block their absorption.

As well as pharmaceutical and veterinary formulations, IgA antibodies produced in accordance with the present invention have applications in the nutritional fields. This may range from the use of the milk *per se* to specific formulations produced containing high
35 IgA levels for well-being, for applications such as nutritional beverages and sports nutrition.

Formulations containing specific IgA against common allergens such as pollens, dust, and

mites for allergy protection are possible. Contemplated herein are formulations containing specific antibodies against mycotoxins, phytotoxins, aflatoxins, pesticides, herbicides, environmental pollutants such as dioxins, polychlorinated biphenyls and fungicides to block absorption of these compounds. Formulations against undesirable food ingredients
5 such as cholesterol to block absorption of these would be particularly useful. In a further aspect the specific IgA may be complexed to probiotics or growth factors for the preparation of formulations for gastric well-being.

In the veterinary corollary, formulations containing IgA for nutritional support particularly
10 of economically important animal offspring such as lambs, piglets, calves, foals and chickens are possible. Formulations consisting of specific IgA directed against undesirable food ingredients such as β carotene to block absorption of these may also be useful.

15 A further area of application for the IgA product of the present invention is in formulations containing the antibodies against skin or hair protein antigens for topical applications, against skin antigens complexed to UV absorbing compounds such as zinc for long-lasting protection against sunburn and with specific IgA antibodies complexed to growth factors for skin repair.

20 The formulations may be prepared in the form of drinks, lotions, powders, creams and the like according to principles well known in the art. The formulations may be for oral, intravenous, intramuscular, subcutaneous, rectal, topical, parenteral administration or such other routes as may be desired.

25 The formulations may include pharmaceutically acceptable carriers or, in the case of nutritional supplements, nutritionally acceptable carriers. Such carriers include aqueous solutions, non-toxic excipients, including salts, preservatives, and buffers. The formulations may include additives such as minerals, vitamins, flavouring agents, scenting
30 agents and the like.

General assistance in the preparation of such formulations may be obtained from Remingtons Pharmaceutical Sciences, 16th Edition. Easton: Mac Publishing Company (1980); the National Formulary XIV, 14th Edition. Washington: American
35 Pharmaceutical Association (1975); and Goodman and Gillmans The Pharmacological basis for Therapeutics (7th Edition), the contents of which are hereby incorporated by reference.

Specific non-limiting examples of the invention will now be described.

EXAMPLE 1

- 5 64 pregnant ewes were selected and divided into eight groups and immunised by intra muscular (IM), intra peritoneal (IP) or intra mammary (IMM) routes or combinations of these three. For the IMM immunisation the right gland only was immunised while the left gland was untreated and acted as the control.

10

11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65
66
67
68
69
70
71
72
73
74
75
76
77
78
79
80
81
82
83
84
85
86
87
88
89
90
91
92
93
94
95
96
97
98
99
100
101
102
103
104
105
106
107
108
109
110
111
112
113
114
115
116
117
118
119
120
121
122
123
124
125
126
127
128
129
130
131
132
133
134
135
136
137
138
139
140
141
142
143
144
145
146
147
148
149
150
151
152
153
154
155
156
157
158
159
160
161
162
163
164
165
166
167
168
169
170
171
172
173
174
175
176
177
178
179
180
181
182
183
184
185
186
187
188
189
190
191
192
193
194
195
196
197
198
199
200
201
202
203
204
205
206
207
208
209
210
211
212
213
214
215
216
217
218
219
220
221
222
223
224
225
226
227
228
229
230
231
232
233
234
235
236
237
238
239
240
241
242
243
244
245
246
247
248
249
250
251
252
253
254
255
256
257
258
259
260
261
262
263
264
265
266
267
268
269
270
271
272
273
274
275
276
277
278
279
280
281
282
283
284
285
286
287
288
289
290
291
292
293
294
295
296
297
298
299
300
301
302
303
304
305
306
307
308
309
310
311
312
313
314
315
316
317
318
319
320
321
322
323
324
325
326
327
328
329
330
331
332
333
334
335
336
337
338
339
340
341
342
343
344
345
346
347
348
349
350
351
352
353
354
355
356
357
358
359
360
361
362
363
364
365
366
367
368
369
370
371
372
373
374
375
376
377
378
379
380
381
382
383
384
385
386
387
388
389
390
391
392
393
394
395
396
397
398
399
400
401
402
403
404
405
406
407
408
409
410
411
412
413
414
415
416
417
418
419
420
421
422
423
424
425
426
427
428
429
430
431
432
433
434
435
436
437
438
439
440
441
442
443
444
445
446
447
448
449
450
451
452
453
454
455
456
457
458
459
460
461
462
463
464
465
466
467
468
469
470
471
472
473
474
475
476
477
478
479
480
481
482
483
484
485
486
487
488
489
490
491
492
493
494
495
496
497
498
499
500
501
502
503
504
505
506
507
508
509
510
511
512
513
514
515
516
517
518
519
520
521
522
523
524
525
526
527
528
529
530
531
532
533
534
535
536
537
538
539
540
541
542
543
544
545
546
547
548
549
550
551
552
553
554
555
556
557
558
559
560
561
562
563
564
565
566
567
568
569
570
571
572
573
574
575
576
577
578
579
580
581
582
583
584
585
586
587
588
589
590
591
592
593
594
595
596
597
598
599
600
601
602
603
604
605
606
607
608
609
610
611
612
613
614
615
616
617
618
619
620
621
622
623
624
625
626
627
628
629
630
631
632
633
634
635
636
637
638
639
640
641
642
643
644
645
646
647
648
649
650
651
652
653
654
655
656
657
658
659
660
661
662
663
664
665
666
667
668
669
670
671
672
673
674
675
676
677
678
679
680
681
682
683
684
685
686
687
688
689
690
691
692
693
694
695
696
697
698
699
700
701
702
703
704
705
706
707
708
709
710
711
712
713
714
715
716
717
718
719
720
721
722
723
724
725
726
727
728
729
730
731
732
733
734
735
736
737
738
739
740
741
742
743
744
745
746
747
748
749
750
751
752
753
754
755
756
757
758
759
760
761
762
763
764
765
766
767
768
769
770
771
772
773
774
775
776
777
778
779
780
781
782
783
784
785
786
787
788
789
790
791
792
793
794
795
796
797
798
799
800
801
802
803
804
805
806
807
808
809
810
811
812
813
814
815
816
817
818
819
820
821
822
823
824
825
826
827
828
829
830
831
832
833
834
835
836
837
838
839
840
841
842
843
844
845
846
847
848
849
850
851
852
853
854
855
856
857
858
859
860
861
862
863
864
865
866
867
868
869
870
871
872
873
874
875
876
877
878
879
880
881
882
883
884
885
886
887
888
889
890
891
892
893
894
895
896
897
898
899
900
901
902
903
904
905
906
907
908
909
910
911
912
913
914
915
916
917
918
919
920
921
922
923
924
925
926
927
928
929
930
931
932
933
934
935
936
937
938
939
940
941
942
943
944
945
946
947
948
949
950
951
952
953
954
955
956
957
958
959
960
961
962
963
964
965
966
967
968
969
970
971
972
973
974
975
976
977
978
979
980
981
982
983
984
985
986
987
988
989
990
991
992
993
994
995
996
997
998
999
1000
1001
1002
1003
1004
1005
1006
1007
1008
1009
1010
1011
1012
1013
1014
1015
1016
1017
1018
1019
1020
1021
1022
1023
1024
1025
1026
1027
1028
1029
1030
1031
1032
1033
1034
1035
1036
1037
1038
1039
1040
1041
1042
1043
1044
1045
1046
1047
1048
1049
1050
1051
1052
1053
1054
1055
1056
1057
1058
1059
1060
1061
1062
1063
1064
1065
1066
1067
1068
1069
1070
1071
1072
1073
1074
1075
1076
1077
1078
1079
1080
1081
1082
1083
1084
1085
1086
1087
1088
1089
1090
1091
1092
1093
1094
1095
1096
1097
1098
1099
1100
1101
1102
1103
1104
1105
1106
1107
1108
1109
1110
1111
1112
1113
1114
1115
1116
1117
1118
1119
1120
1121
1122
1123
1124
1125
1126
1127
1128
1129
1130
1131
1132
1133
1134
1135
1136
1137
1138
1139
1140
1141
1142
1143
1144
1145
1146
1147
1148
1149
1150
1151
1152
1153
1154
1155
1156
1157
1158
1159
1160
1161
1162
1163
1164
1165
1166
1167
1168
1169
1170
1171
1172
1173
1174
1175
1176
1177
1178
1179
1180
1181
1182
1183
1184
1185
1186
1187
1188
1189
1190
1191
1192
1193
1194
1195
1196
1197
1198
1199
1200
1201
1202
1203
1204
1205
1206
1207
1208
1209
1210
1211
1212
1213
1214
1215
1216
1217
1218
1219
1220
1221
1222
1223
1224
1225
1226
1227
1228
1229
1230
1231
1232
1233
1234
1235
1236
1237
1238
1239
1240
1241
1242
1243
1244
1245
1246
1247
1248
1249
1250
1251
1252
1253
1254
1255
1256
1257
1258
1259
1260
1261
1262
1263
1264
1265
1266
1267
1268
1269
1270
1271
1272
1273
1274
1275
1276
1277
1278
1279
1280
1281
1282
1283
1284
1285
1286
1287
1288
1289
1290
1291
1292
1293
1294
1295
1296
1297
1298
1299
1300
1301
1302
1303
1304
1305
1306
1307
1308
1309
1310
1311
1312
1313
1314
1315
1316
1317
1318
1319
1320
1321
1322
1323
1324
1325
1326
1327
1328
1329
1330
1331
1332
1333
1334
1335
1336
1337
1338
1339
1340
1341
1342
1343
1344
1345
1346
1347
1348
1349
1350
1351
1352
1353
1354
1355
1356
1357
1358
1359
1360
1361
1362
1363
1364
1365
1366
1367
1368
1369
1370
1371
1372
1373
1374
1375
1376
1377
1378
1379
1380
1381
1382
1383
1384
1385
1386
1387
1388
1389
1390
1391
1392
1393
1394
1395
1396
1397
1398
1399
1400
1401
1402
1403
1404
1405
1406
1407
1408
1409
1410
1411
1412
1413
1414
1415
1416
1417
1418
1419
1420
1421
1422
1423
1424
1425
1426
1427
1428
1429
1430
1431
1432
1433
1434
1435
1436
1437
1438
1439
1440
1441
1442
1443
1444
1445
1446
1447
1448
1449
1450
1451
1452
1453
1454
1455
1456
1457
1458
1459
1460
1461
1462
1463
1464
1465
1466
1467
1468
1469
1470
1471
1472
1473
1474
1475
1476
1477
1478
1479
1480
1481
1482
1483
1484
1485
1486
1487
1488
1489
1490
1491
1492
1493
1494
1495
1496
1497
1498
1499
1500
1501
1502
1503
1504
1505
1506
1507
1508
1509
1510
1511
1512
1513
1514
1515
1516
1517
1518
1519
1520
1521
1522
1523
1524
1525
1526
1527
1528
1529
1530
1531
1532
1533
1534
1535
1536
1537
1538
1539
1540
1541
1542
1543
1544
1545
1546
1547
1548
1549
1550
1551
1552
1553
1554
1555
1556
1557
1558
1559
1560
1561
1562
1563
1564
1565
1566
1567
1568
1569
1570
1571
1572
1573
1574
1575
1576
1577
1578
1579
1580
1581
1582
1583
1584
1585
1586
1587
1588
1589
1590
1591
1592
1593
1594
1595
1596
1597
1598
1599
1600
1601
1602
1603
1604
1605
1606
1607
1608
1609
1610
1611
1612
1613
1614
1615
1616
1617
1618
1619
1620
1621
1622
1623
1624
1625
1626
1627
1628
1629
1630
1631
1632
1633
1634
1635
1636
1637
1638
1639
1640
1641
1642
1643
1644
1645
1646
1647
1648
1649
1650
1651
1652
1653
1654
1655
1656
1657
1658
1659
1660
1661
1662
1663
1664
1665
1666
1667
1668
1669
1670
1671
1672
1673
1674
1675
1676
1677
1678
1679
1680
1681
1682
1683
1684
1685
1686
1687
1688
1689
1690
1691
1692
1693
1694
1695
1696
1697
1698
1699
1700
1701
1702
1703
1704
1705
1706
1707
1708
1709
1710
1711
1712
1713
1714
1715
1716
1717
1718
1719
1720
1721
1722
1723
1724
1725
1726
1727
1728
1729
1730
1731
1732
1733
1734
1735
1736
1737
1738
1739
1740
1741
1742
1743
1744
1745
1746
1747
1748
1749
1750
1751
1752
1753
1754
1755
1756
1757
1758
1759
1760
1761
1762
1763
1764
1765
1766
1767
1768
1769
1770
1771
1772
1773
1774
1775
1776
1777
1778
1779
1780
1781
1782
1783
1784
1785
1786
1787
1788
1789
1790
1791
1792
1793
1794
1795
1796
1797
1798
1799
1800
1801
1802
1803
1804
1805
1806
1807
1808
1809
1810
1811
1812
1813
1814
1815
1816
1817
1818
1819
1820
1821
1822
1823
1824
1825
1826
1827
1828
1829
1830
1831
1832
1833
1834
1835
1836
1837
1838
1839
1840
1841
1842
1843
1844
1845
1846
1847
1848
1849
1850
1851
1852
1853
1854
1855
1856
1857
1858
1859
1860
1861
1862
1863
1864
1865
1866
1867
1868
1869
1870
1871
1872
1873
1874
1875
1876
1877
1878
1879
1880
1881
1882
1883
1884
1885
1886
1887
1888
1889
1890
1891
1892
1893
1894
1895
1896
1897
1898
1899
1900
1901
1902
1903
1904
1905
1906
1907
1908
1909
1910
1911
1912
1913
1914
1915
1916
1917
1918
1919
1920
1921
1922
1923
1924
1925
1926
1927
1928
1929
1930
1931
1932
1933
1934
1935
1936
1937
1938
1939
1940
1941
1942
1943
1944
1945
1946
1947
1948
1949
1950
1951
1952
1953
1954
1955
1956
1957
1958
1959
1960
1961
1962
1963
1964
1965
1966
1967
1968
1969
1970
1971
1972
1973
1974
1975
1976
1977
1978
1979
1980
1981
1982
1983
1984
1985
1986
1987
1988
1989
1990
1991
1992
1993
1994
1995
1996
1997
1998
1999
2000
2001
2002
2003
2004
2005
2006
2007
2008
2009
2010
2011
2012
2013
2014
2015
2016
2017
2018
2019
2020
2021
2022
2023
2024
2025
2026
2027
2028
2029
2030
2031
2032
2033
2034
2035
2036
2037
2038
2039
2040
2041
2042
2043
2044
2045
2046
2047
2048
2049
2050
2051
2052
2053
2054
2055
2056
2057
2058
2059
2060
2061
2062
2063
2064
2065
2066
2067
2068
2069
2070
2071
2072
2073
2074
2075
2076
2077
2078
2079
2080
2081
2082
2083
2084
2085
2086
2087
2088
2089
2090
2091
2092
2093
2094
2095
2096
2097
2098
2099
2100
2101
2102
2103
2104
2105
2106
2107
2108
2109
2110
2111
2112
2113
2114
2115
2116
2117
2118
2119
2120
2121
2122
2123
2124
2125
2126
2127
2128
2129
2130
2131
2132
2133
2134
2135
2136
2137
2138
2139
2140
2141
2142
2143
2144
2145
2146
2147
2148
2149
2150
2151
2152
2153
2154
2155
2156
2157
2158
2159
2160
2161
2162
2163
2164
2165
2166
2167
2168
2169
2170
2171
2172
2173
2174
2175
2176
2177
2178
2179
2180
2181
2182
2183
2184
2185
2186
2187
2188
2189
2190
2191
2192
2193
219

Immunisation Protocol

	Group (n = 8)	Immunisation Route
5	1	IM
	2	IP
	3	IMM
	4	IM/IP
	5	IM/IMM
10	6	IP/IMM
	7	IM/IP/IMM
	8	NIL (Control)

- 15 Animals were immunised according to the following schedule.

Immunisation Schedule

	I	II	III	IV		V	VI	VII
20	-8 wk	-4 wk	-2 wk	-1 wk	0	+4wk	+8 wk	+12 wk
					Parturition			Weaning
					n			
25	IM-1	IM-2		IM-3		IM-4	IM-5	IM-6
	IP-1	IP-2		IP-3		IP-4	IP-5	IP-6
		IMM-1	IMM-2					IMM-3

Antigen

- 30 A commercially available pathogenic *E. coli* vaccine (Suvaxyn, Maternafend-4; J&H Pacific Ltd, NZ) was used as antigen. This vaccine is known to have high concentrations of K88, K99, 987P and F41 pili antigens.

- 35 Details of the immunisation protocols are as follows:

*Immunisation I.**IM/IP*

- 40 Stock vaccine was emulsified with Freund's Incomplete Adjuvant (FIC; 1 part vaccine: 3 parts FIC).

IM; 1 ml per site; 2 sites

IP; 1 ml per site; 1 site

Immunisation II.

5 IM/IP

Repeat IM/IP as for Immunisation I.

IMM

Stock vaccine was diluted in sterile saline (1 part vaccine: 1 part saline). Antibiotic
10 (Dupocillin) was added to the IMM immunogen in the ratio of 1ml antibiotic: 30 ml immunogen.

IMM; 1 ml per gland; right gland only

Immunisation III

15

IMM

Repeat Immunisation II for IMM only.

Immunisation IV, V and VI

20

IM/IP

Repeat Immunisation I

Immunisation VII

25

IM/IP & IMM

Repeat Immunisation II

30 *Animal Health Status*

The general health of the ewes in the trial was monitored by regular weight checks and veterinary inspections. There was no discernible difference observed in weight gain/loss between the treatment groups. No adverse effects of immunisation were observed. Two
35 of the 64 ewes were treated for mastitis, one ewe from Group 6 (IP/IMM) and one ewe from Group 7 (IM/IP/IMM). Overall, no deleterious effects of IMM immunisation were noted.

Samples

The ewes were bled before Immunisation I, II, and IV, prior to lambing. *Post parturition* samples of blood and colostrum/milk (left and right mammary glands separately) were collected at Day 0 (parturition), 1, 2, 3, and 5, then Week 1, 2 and 3, then Month 1, 2 and 3.

Bloods were collected on ice into EDTA vacutainers. Separated plasma was stored at -20 °C for antibody analysis. Left (untreated) and right (immunised) mammary gland colostrum/milk samples (20-30 ml) were kept on ice until centrifuged (4°C; 20 minutes; 2,000 g_{max}) to remove fat. Skimmed supernatant was re-centrifuged (4°C; 1 hour; 40,000 g_{max}) to separate milk whey/plasma and casein. Supernatant was stored at -20 °C for antibody analysis.

15 Sample analysis

All samples and reagents were diluted with 0.01M phosphate buffered saline (pH 7.5) containing 0.05% v/v Tween-20 (PBS-T) and 1% w/v Bovine Serum Albumin (BSA; type A7030, Sigma Chem. Co., USA) and all washes were carried out by an automated plate washer (ELP-35, BioTek Instruments, USA) using PBS-T, unless otherwise stated.

- ELISA plates (Maxisorp F-96 immunoplates, Nunc, Denmark) were coated with 100µl of *E. coli* antigen (Suvaxyn Maternafend-4; J & H Pacific Ltd, NZ) diluted 1:1K in 0.05M carbonate buffer (pH 9.6), incubated overnight at 4°C and washed three times. Remaining activated sites on immunoplates were blocked by incubating 2 hours at 22°C with 250µl
- 5 PBS-T containing 1% w/v BSA. After washing plates 2 times, 100µl of 10-fold serial dilutions of test samples (primary antibody; 1:100, 1:1K, 1:10K, 1:100K) were added to duplicate wells. Plates were incubated 2 hours at 22°C then washed 3 times. 100µl of second antibody consisting of heavy-chain specific rabbit anti-sheep IgA (1:200K; Bethyl Laboratories, USA) were added to the plates. Plates were incubated overnight at 4°C,
- 10 then washed 3 times prior to the addition of 100µl of the enzyme conjugate, goat anti-rabbit Ig conjugated to horse radish peroxidase (1:8K; Dako, Denmark). After a 2 hour incubation at 22°C, the plates were washed 2 times with PBS-T then 2 times with PBS containing no Tween-20 and filled with 100µl of freshly prepared substrate solution. The substrate solution consisted of 0.1 g/l 3,3',5,5'-tetramethylbenzidine (Boehringer
- 15 Mannheim, Germany) in 0.1M sodium acetate buffer (pH 5.5) containing 1.3 mmol/l hydrogen peroxide. Following a 30 minute incubation at 22°C, 50µl of stopping solution, 2M H₂SO₄, were added and the optical density (OD) was measured at 450nm by an automated plate reader (EL311s, BioTek Instruments, USA).
- 20 With each ELISA microplate a positive quality control sample (assayed at 1:100, 1:1K, 1:10K, 1:100K and 1:1000K) and a negative quality control sample (assayed at 1:1K) was run with the samples. Absorbance values from these control samples were used in calculations to determine sample antibody titres. The median absorbance between the maximum absorbance of the positive control and the absorbance of the negative control
- 25 gives a 50 % figure. The reciprocal dilution of sample antibody equivalent to this 50 % absorbance figure is classified as the antibody titre for the sample.

Figure 1 shows a typical dilution curve for the positive control sample in the *E. coli* IgA ELISA.

30

Results

Samples were initially assayed as group pools to obtain an overview of the group responses to the different immunisation regimens.

35

Figure 2 shows the right (immunised) gland milk anti-*E. coli* IgA titres for all groups on Day 0, Day 5, Week 2 and Week 4. In all groups, IgA milk antibody titres were highest on Day 0 (parturition) with levels falling over the first week. Group 7 (IM/IP/IMM) gave

the best IgA response with a milk antibody titre of 105K. This was followed by Group 5 (IM/IMM) with a titre of 50K and Group 6 (IP/IMM) with a titre of 22K. The other groups gave a minimal response including Group 3 (IMM). By Day 5, the IgA milk antibody titres in Groups 5, 6 and 7 had fallen to about 20% of Day 0 titres but by Week 4 the titres were still significant being approximately 3K.

Figure 3 contrasts anti-*E. coli* IgA responses of the right (immunised) gland and the left (untreated) gland on Day 0. IgA milk antibody titres showed a marked difference in response between samples from the right (immunised) gland and the left (untreated) gland. While high titres were measured in the right milk samples of Groups 5, 6 and 7, the titres in the corresponding left milk samples were, at best, only 20% of these levels.

EXAMPLE 2

Individual samples analysis of anti-*E. coli* IgA titre responses was conducted for each of the immunisation groups of Example 1. Samples were analysed using the ELISA assay according to the process of Example 1. The results for Day 0 and 1 are shown in Figures 4 and 5, respectively.

Results

In general agreement with the earlier pooled data, titres were low in animals immunised by IM, IP or IMM routes alone and much higher in animals treated by the combination IM/IMM, IP/IMM and IM/IP/IMM routes (Mean \pm s.e.m. antibody titres; 1/32,100 \pm 1/13,500; 1/29,000 \pm 1/12,000; 1/30,000 \pm 1/7,300, respectively).

No significant differences were observed in mean IgA titre responses resulting from immunisation by each of the three combination routes. However, substantial within-group variability in titre response was observed and notably the standard error of the mean for the IM/IP/IMM group (1/7,300) was much lower than that calculated for the IM/IMM (1/13,500) and IP/IMM (1/12,500) routes.

This pattern of titre response was maintained in milk samples from Day 2 and subsequently. The data appear to indicate that immunisation by the three site IM/IP/IMM procedure does not increase the magnitude of the response above those obtained with the IM/IMM and IP/IMM combinations but serves to decrease the between animal variability in IgA response.

EXAMPLE 3

64 pregnant ewes were selected and divided into eight groups and immunised by intra muscular (IM), intra peritoneal (IP) or intra mammary (IMM) routes or combinations of these three. For the IMM immunisation the right gland only was immunised while the left gland was untreated and acted as the control.

Immunisation Protocol

	Group (n = 8)	Immunisation Route
10	1	IM
	2	IP
	3	IMM
15	4	IM/IP
	5	IM/IMM
	6	IP/IMM
	7	IM/IP/IMM
20	8	NIL - Control

Animals were immunised according to the following schedule.

Immunisation Schedule

	I -8 wk	II -4 wk	III -2 wk	IV -1 wk	0	V +4wk	VI +8 wk	VII +12 wk
	↓	↓	↓	↓		↓	↓	↓
					Parturition			Weaning
					n			
30	IM-1 IP-1	IM-2 IP-2 IMM-1	IMM-2	IM-3 IP-3		IM-4 IP-4	IM-5 IP-5	IM-6 IP-6 IMM-3

35 *Antigen*

A commercially available vaccine, 3K Scourguard (SmithKline Beecham, Royal Oak, Auckland, New Zealand) was used as immunogen. The vaccine contains pathogenic *E. coli*, bovine rotavirus and coronavirus.

40

Details of the immunisation protocols are as follows:

The immunisation protocol of Example 1 was repeated with 3K Scourguard used as stock vaccine in place of Maternafend.

Animal Health Status

5

The general health of the ewes in the trial was monitored by regular weight checks and veterinary inspections. And as for Example 1, there was no discernible difference observed in weight gain/loss between the treatment groups and no adverse effects of immunisation were observed.

10

Samples

Samples of blood and colostrum/milk were taken according to the protocol of Example 1.

15

Sample analysis

The ELISA assay was performed according to the method of Example 1, with the exception that 3K Scourguard (1:1K) was used for microplate coating in place of Maternafend. Blood plasma and colostrum/milk were pooled to obtain an initial indication of group antibody responses.

20

Results

Figure 6 shows the right (immunised) gland milk anti-3K Scourguard IgA titres for all groups on Day 0 and 5. In all groups, IgA milk antibody titres were highest on Day 0 (parturition) with levels falling over the first week. Group 7 (IM/IP/IMM) gave the best IgA response with a milk antibody titre of 210K. This was followed by Group 5 (IM/IMM) with a titre of 70K and Group 3 with a titre of 27K. Group 6 (IP/IMM) gave a milk antibody response of 20K. The other groups gave a minimal response. By Day 5, the IgA milk antibody titres in Group 3, 5 and 7 had fallen to about 10 % of Day 0 titres.

30

Figure 7 shows the right (immunised) and left (untreated) gland milk anti-3K Scourguard IgA titres for all groups on Day 0. The right gland had a much greater response than that of the left gland. There was no significant antibody titre response for the untreated left gland except for Group 3.

35

EXAMPLE 4

32 pregnant ewes were assigned to four treatment groups and immunised by combinations of intra muscular (IM), intra peritoneal (IP) or intra mammary (IMM) routes. IMM immunogen for Group 3 was in aqueous solution while IMM immunogen for Group 4 was emulsified in FIC. For the IMM immunisations the right gland only was immunised while the left gland was untreated and acted as the control.

Immunisation Protocol

10

Group (n = 8)	Immunisation Route
1	IM/IMM
2	IP/IMM
3	IM/IP/IMM
4	IM/IP/IMM(FIC)

15

Animals were immunised according to the following schedule.

20

Immunisation Schedule

I -8 wk	II -4 wk	III -2 wk	IV -1 wk	0
↓	↓	↓	↓	
IM-1 IP-1	IMM-1 IMM(FIC)-1 IM-2 IP-2	IMM-2 IMM(FIC)-2	IM-3 IP-3	Parturition

25

30

Antigen

A TNF preparation, commercially available from R & D Systems, 614 McKinley Place, New England, USA, was used as antigen. A stock solution (1mg/ml) was prepared by reconstituting the freeze dried TNF in sterile saline.

35

Details of the immunisation protocols are as follows:

40

Immunisation I.

IM/IP

Stock antigen solution was diluted to 0.16 mg/ml then emulsified with FIC (1 part saline:

5 3 parts FIC).

IM; 1 ml per site; 2 sites

IP; 1 ml per site; 1 site

Immunisation II.

10

IM/IP

Repeat IM/IP as for Immunisation I.

IMM

15 For right glands: Stock antigen solution was diluted to 0.1 mg/ml in sterile saline. Antibiotic (Dupocillin) was added in the ratio of 1 ml antibiotic: 40 ml immunogen.

IMM; 1 ml per right gland

IMM(FIC)

20 For right glands: Stock antigen solution was diluted to 0.32 mg/ml in sterile saline and emulsified with FIC (1 part saline: 3 parts FIC). Antibiotic (Dupocillin) was added in the ratio of 1 ml antibiotic: 40 ml immunogen.

IMM (FIC); 1 ml per right gland

25 *Immunisation III*

IMM/IMM(FIC)

Repeat Immunisation II for IMM/IMM(FIC) only.

30 *Immunisation IV*

IM/IP

Repeat Immunisation I

35 *Animal Health Status*

The general health of the ewes in the trial was monitored by regular weight checks and veterinary inspections. The animals maintained weight during pregnancy and lactation

and no between group effects of treatment were observed. Two animals died from unrelated causes during pregnancy/lambing (both ewes from Group 1, IM/IMM). One animal was withdrawn from the trial due to mastitis in the left untreated gland. No adverse effects on immunisation were noted. Evidence of ulceration at IM and IP immunisation sites was minimal. No significant differences were observed in mammary function between the left and right glands and between glands immunised with immunogen in sterile saline or FIC.

Samples

10

The ewes were bled before Immunisation I, II, and IV, prior to lambing. Post lambing, samples of blood and colostrum/milk (left and right mammary glands separately) were collected at Day 1 (parturition), 2, 3, 6, 14 and 28, and Month 2 and 3.

15 Samples were collected and treated according the format used in Example 1.

Sample Analysis

20 The ELISA assay for TNF was performed according to the format used for Example 1, with the exception that TNF was used for plate coating (2 mg/ml). Figure 8 shows a typical dilution curve for the positive and negative control samples in the TNF IgA ELISA. Inter assay precision was calculated from 10 repeat analysis of the positive control and the coefficient of variation was 10.2 %.

Results

IgA milk responses

Individual sample analyses of anti-TNF IgA titre responses in the right (immunised) gland Day 1 *post-partum* milk samples for each of the immunisation groups are shown in Figure 9. Titres were low in animals immunised by the IM/IMM, IP/IMM or IM/IP/IMM routes where the immunogen was administered in saline solution. (Mean \pm s.e.m. antibody titres: $1/3,600 \pm 1/2,600$; $1/3,300 \pm 1/1,500$; $1/900 \pm 1/500$, respectively.) By contrast, titres were some 20-fold higher in animals treated by the combination IM/IP/IMM routes where the IMM immunogen was emulsified in FIC ($1/61,900 \pm 1/29,600$). Considerable variation was observed in the responses of the individual Group 4 animals with titres ranging from $1/4,000$ to $1/250,000$.

Right and Left Gland Milk Antibody Responses

Significant differences were seen in anti-TNF IgA milk titres from right (immunised) and left (untreated) glands with the left gland IgA titre being almost undetectable, in agreement with earlier findings for *E. coli*. Figure 10 depicts the data for anti-TNF IgA titres for Day 1 *post-partum* milk samples from the left and right glands.

Milk Antibody Response Over Lactation

The relationship of anti-TNF IgA to stage of lactation is shown in Figure 11. Data are mean \pm s.e.m. right (immunised) gland milk titres from Group 4 animals treated by the combination IM/IP/IMM(FIC) route. Anti-TNF IgA titres were highest in initial *post-partum* mammary secretions and found to decline to approximately 10% of peak levels by Day 6 and approximately 5 % by Month 1 (equivalent to an IgA titre of 1/3,500). The overall pattern of response was similar to that seen in the *E. coli* trial, the antibody decline coinciding with the onset of full lactation and increasing milk volumes.

EXAMPLE 5

35 pregnant cows were divided into four groups and immunised by either two routes, three routes or none according to the protocol below. Immunogen was emulsified in Freund's Incomplete Adjuvant (FIC) for right sided glands intra mammary (IMM) immunisations and for intra muscular (IM) and intra peritoneal (IP) immunisations. Left sided glands utilised immunogen in aqueous solution.

Immunisation Protocol

Group	Immunisation Route
1a (n = 10)	IM/IMM
1b (n = 10)	IP/IMM
1c (n = 10)	IM/IP/IMM
1d (n = 5)	NIL

Animals were immunised according to the following schedule.

Immunisation schedule

	I	II	III	IV	
	-8 wk	-4 wk	-2 wk	-1 wk	0
5					
	IM-1	IMM-1	IMM-2		Parturition
	IP-1	IM-2		IM-3	
10		IP-2		IP-3	

Antigen

Antigen for immunisation was the yeast, *Candida albicans*. The yeast cells were cultured in medium, harvested by centrifugation, washed and heat killed then freeze dried. A stock solution of *C. albicans* (7 mg protein per ml) was prepared by reconstituting the freeze dried *C. albicans* in phosphate buffer.

Details of the immunisation protocols are as follows:

*Immunisation I.**IM/IP*

Stock antigen solution was diluted to 1mg/ml in sterile saline and emulsified with FIC (1 part saline: 3 parts FIC).

IM; 2 ml per site; 2 sites

IP; 4 ml per site; 1 site

*Immunisation II.**IM/IP*

Repeat IM/IP as for Immunisation I.

IMM

For right glands: Stock antigen solution was diluted to 1mg/ml in sterile saline and emulsified with FIC (1 part saline: 3 parts FIC).

For left glands: Stock antigen solution was diluted to 0.25mg/ml in sterile saline.

IMM; 2 ml per gland; 4 glands

Immunisation III

IMM

Repeat Immunisation II for IMM only.

Immunisation IV

5

IM/IP

Repeat Immunisation I

Animal Health Status

10

The general health of the cows in the trial was monitored by regular weight checks and veterinary inspections. Immunisation sites were inspected at regular intervals to assess effects of the immunisation procedure. No clinical grade site reactions were observed at any of the sites immunised. In addition, milk volume data collected indicated that treatment of the mammary gland did not effect the overall lactation performance of the animals.

15

Samples

20 The cows were bled before Immunisation I, II, and IV, prior to calving. *Post parturition* samples of blood and colostrum/milk were collected at Day 1, 2, 3, 5, 7, 14, 28 and 60. On sample days, cows were quarter milked (ie. samples were collected from individual glands) AM and PM, milk volumes were recorded and 100ml sample retained. AM and PM quarter milk samples were pooled for laboratory analyses.

25

Blood and colostrum/milk samples were treated according to the format used for Example 1.

Sample Analysis

30

The ELISA assay for *C albicans* was performed according to the format used for Example 1, with the exception that: *C albicans* (5mg/ml) was used for plate coating; second antibody used to identify class specificity was rabbit anti-bovine IgA (1:40K; Bethyl Laboratories, USA); enzyme conjugated antibody was goat anti-rabbit (1:12K; Dako, Denmark). The end point detection system was the same as in Example 1.

35

Figure 12 shows a typical antibody dilution curve for the positive control sample.

Results

For Groups 1a, 1b and 1c, the anti-*C. albicans* IgA milk antibody titres were highest on Day 1 (parturition) with levels falling over the lactation period, as observed with the earlier sheep trials (Examples 1-4).

Two site versus three site immunisation

Figure 13 shows the right (IMM immunogen in FIC) gland milk anti-*C. albicans* IgA antibody titres for all groups on Day 1, 2, 7, 14 and 60. Animals immunised at three sites (Group 1c; IM/IP/IMM) had a much higher response than animals immunised at two sites (Group 1a; IM/IMM and Group 1b; IP/IMM). The mean \pm s.e.m. antibody titres were $11,700 \pm 3,700$, $2,500 \pm 700$ and $3,000 \pm 900$, respectively. Group 1d (Control) had a very low antibody titre on Day 1 (titre of 200 ± 70). The higher titres for animals immunised at three sites was maintained over the lactation period.

Freund's versus saline for IMM immunisations

Figure 14 shows the milk anti-*C. albicans* IgA titre comparison of the left (aqueous IMM immunogen) and right (FIC IMM immunogen) glands in Group 1c over the lactation period. Glands immunised with immunogen emulsified in FIC gave a higher response than glands immunised with aqueous immunogen and this difference increased with time. On Day 1 the right gland titre was $11,700 \pm 3,700$ compared to the left gland titre of $8,700 \pm 2,900$. By Day 60, the left gland titre had declined to 590 ± 200 while the right gland titre was 4 fold higher ($2,300 \pm 800$). This difference between left and right gland response was also apparent in the groups immunised at two sites (Group 1a and 1b).

Thus, in accordance with the present invention there is provided a process for the induction of IgA in a mammal, and the production of IgA in mammalian milk at levels higher than have previously been obtained or might have been anticipated from combining a third administration route for an antigen with known two route administration protocols. Alternatively, the present invention at least provides a method whereby the between animal variability in IgA antibody titre response can be reduced. It will be appreciated that these results represent an advantage where products containing IgA are sought for use in pharmaceutical, veterinary and cosmetic formulations as well as in nutritional and dietary supplements.

It will be further appreciated by those persons skilled in the art that the present description

is provided by way of example only and that the scope of the invention is not limited thereto.

REFERENCES

- 5 1. Immune Response of Pregnant cows to Bovine Rotavirus Immunisation, Saif L. J. *et al*; American Journal of Veterinary Research (1984), 45: 1, 49 - 58.
2. Comparative Effect of Selected Adjuvants on the Response in the Bovine Mammary Gland to Staphylococcal and Streptococcal antigens, Opdebeeck, J. P. and Norcross, N. L; Veterinary Immunopathology (1984), 6: 341-351.
- 10 3. Novel Vaccination Strategies for the Control of Mucosal Infection, Husband, A. J; Vaccine (1993), 11: 2, 107-112.
4. Immunological Effector Mechanisms in Ruminants, D.L. Watson, I. G. Colditz, H. S. Gill; in Vaccines in Agriculture (1994), pp 21-36, eds Wood *et al*, CSIRO, Australia.
- 15 5. Responses of Antibody Titers to Intramammary Immunization with *Escherichia coli* J5 Bacterin; J. S. Hogan, K. L. Smith, P. Schoenberger, S. Romig and L. Thompson; J Dairy Sci. (1997) 80: 2398-2402.
6. Immune Mechanisms of the Ruminant Mammary Gland, R. F. Sheldrake; Australian Journal of Dairy Technology (1987), 42:1-2, 30-32.
- 20 7. The Effect of Intraperitoneal and Intramammary immunization of sheep on the numbers of antibody-containing cells in the mammary gland, and antibody titres in blood serum and mammary secretions; R. F. Sheldrake, A. J. Husband, D. L. Watson, A. W. Cripps; Immunology (1985), 56: 605-614.
8. Specific antibody-containing cells in the mammary gland of non-lactating sheep after intraperitoneal and intramammary immunisation; R. F. Sheldrake, A. J. Husband, D. L. Watson, Research in Veterinary Science (1985), 38: 312-316.
- 25 9. Vaccination against enteric Rota and Coronaviruses in cattle and pigs: enhancement of lactogenic immunity, C. F. Crouch; Vaccine (1985), 3: 284-291.
10. The mucosal immune system with particular reference to ruminant animals, A. K. Lascelles, K. J. Beh, T. K. Mukkur, D. L. Watson; in The Ruminant Immune System in Health and Disease (1986), pp 429-457, eds Morrison, Cambridge University Press, UK
- 30 11. Origin of antibody-containing cells in the ovine mammary gland following intraperitoneal and intramammary immunisation, R. F. Sheldrake, A. J. Husband, D. L. Watson; Research in Veterinary Science (1988), 45, 156-159.
- 35 12. IgA immune responses in the respiratory tract of pigs; R. F. Sheldrake; Research in Veterinary Science, (1990), 49, 98-103.
13. The Development of Vaccines to Protect Mucosal Surfaces; A. J. Husband, M. L. Dunkley, V. L. Clifton, in Health and Production for the 21st Century (1993), pp

- 82-88, ed Beh, CSIRO, Australia
14. Treatment of Infantile *E. coli* Gastroenteritis with Specific Bovine anti-*E. coli* Milk Immunoglobulins; C. Mietens, H. Keinhorst, H. Hilpert, H. Gerber, H. Amster, J.J. Pahud; Eur. J. Pediatr. (1979), 132: 239-252.
 - 5 15. Bovine Milk Antibodies in the Treatment of Enteric Infections and their ability to eliminate virulence factors from pathogenic *E. coli*; J. J. Pahud, H. Hilpert, K. Schwarz, H. Amster, M. Smiley; in Advances in Exp. Med & Biol.: The Ruminant Immune System (1981), pp 591-600, ed Butler, Plenum Press, NY
 16. Use of Bovine Milk Concentrate Containing Antibody to Rotavirus to treat
10 Rotavirus Gastroenteritis in Infants; H. Hilpert, H. Brussow, C. Mietens, J. Sidoti, L. Lerner, H. Werchau; Journal of Infectious Diseases (1987), 156: 1, 158-166.
 17. Bovine Milk Immunoglobulins for Passive Immunity to Infantile Rotavirus Gastroenteritis; H. Brussow, H. Hilpert, I. Walther, J. Sidoti, C. Mietens, P. Bachmann; Journal of Clinical Microbiology (1987), 25: 6, 982-986.
 - 15 18. Antibodies: A Laboratory Manual (1988), p110-114; ed Harlan Cold Spring Water Laboratory, NY.
 19. Review of Polyclonal Antibody Production Procedures, Hanly et al; ILAR Journal (1995), 37:3, 93-118.
 20. Purification of Secretory IgA from Bovine Colostrum, Kanamaru et al; Agric. Biol. Chem (1982), 46:16, 1531-1537.
 21. A Method for Preparation of IgA from Bovine Mammary Secretions, Nielson, K; Can. J. Vet. Res (1986); 50: 227-231.
 22. Ultrafiltration and Gel Filtration methods for separation of Immunoglobulins with secretory component from bovine milk, Kanamaru et al; Milchwissenschaft (1993),
25 48: 5, 247-251.
 23. Antibodies from Milk for the prevention and Treatment of Diarrheal disease, Ruiz, L.P; in Indigenous Antimicrobial Agents of Milk - Recent Developments (1994), pp 108-121, International Dairy Federation, Brussels, Belgium.
 24. Chronic Cryptosporidial Diarrhoea and Hyperimmune Cow Colostrum, Tzipori, Roberton, Cooper, White; Lancet (1987), II: 8554, 344-345.
 - 30 25. Production of Hyperimmune Bovine Colostrum against *Campylobacter* Jejuni; Husu, Syvaioja, Ahola-Luttila, Kalsta, Sivela, Kosunen; Journal of Applied Bacteriology (1993), 74:5, 564-559.
 26. Production and Preparation of Hyperimmune Bovine Colostrum for passive
35 Immunotherapy of Cryptosporidiosis; Fayer, Tilley, Upton, Guidry, Thayer, Hildreth, Thomson; Journal of Protozoology (1991), 38:6, 38S-39S
 27. Efficacy of Bovine Milk Immunoglobulin Concentrate in Preventing Illness after *Shigella* Flexneri Challenge; Tacket, Binion, Bostwick, Losonsky, Roy, Edelman;

- American Journal of Tropical Medicine and Hygiene, (1992) 47:3, 276-283.
28. Protection of Gnotobiotic rats against dental caries by passive immunization with Bovine Milk Antibodies to Streptococcus mutans; Michalek, Gregory, Harmon, Katz, Richardson, Hilton, Filler, McGhee; Infection and Immunity (1987), 55: 10, 2341-2347.
- 5 29. Cessation of Cryptosporidium-associated Diarrhea in an acquired Immunodeficiency Syndrome Patient after Treatment with Hyperimmune Bovine Colostrum; Ungar, Ward, Fayer, Quinn; Gastroenterology New York (1990), 90: 2, 486-489.
- 10 30. Bovine Colostrum Immunoglobulin Concentrate for Cryptosporidiosis in AIDS; Heaton; in Archives of Disease in Childhood (1994), 70: 4, 356-357.

15 All articles and patents referenced here and in the disclosure are incorporated by reference.